

# Production protocol

## Lentivirus

### Safe Use of Lentivirus (Lv)

---

1. Lentivirus (Lv) related experiments should be conducted in biosafety level 2 facilities (BL-2 level).
2. Please equip with lab coat, mask, gloves completely, and try your best to avoid exposing hand and arm.
3. Be careful of splashing virus suspension. If biosafety cabinet is contaminated with virus during operation, scrub the table-board with solution comprising 70% alcohol and 1% SDS immediately. All tips, tubes, culture plates, medium contacting virus must be soaked in chlorine-containing disinfectant before disposal.
4. If centrifuging is required, a centrifuge tube should be tightly sealed. Seal the tube with parafilm before centrifuging if condition allowed.
5. Lentivirus related animal experiments should also be conducted in BL-2 level.
6. Lentivirus associated waste materials need to be specially collected and autoclaved before disposal.
7. Wash hands with sanitizer after experiment.

### Storage and Dilution of Lentivirus

---

#### Storage of Lentivirus

Virus can be stored at 4°C for a short time (less than a week) before using after reception. Since Lentiviruses are sensitive to freeze-thawing and the titer drops with repeated freeze-thawing, aliquot viral stock should be stored at -80°C freezer immediately upon arrival for long-term usage. While virus titer redetection is suggested before using if the lentiviruses have been stored for more than 12 months.

#### Dilution of Lentivirus

Dissolve virus in ice water if virus dilution is required. After dissolving, mix the virus with medium, sterile PBS or normal saline solution, keeping at 4°C (using within a week).

#### *Precautions*

- Avoid lentivirus exposure to environmental extremes (pH, chelating agents like EDTA, temperature, organic solvents, protein denaturants, strong detergents, etc.)
- Avoid introducing air into the lentivirus samples during vortex, blowing bubbles or similar operations, which may result in protein denaturation.

- Avoid repeated freezing and thawing.
- Avoid exposing to “regular” plastics (especially polystyrene or hydrophobic plastics) for prolonged periods in liquid phase. Most lentivirus viruses are very sticky and loss can occur if exposed to regular plastics, including tubes, cell culture plates, pipette tips, if not frozen. It is best to store lentivirus in siliconized or low protein binding tubes. Pluronic F-68 used at 0.01%-0.1% in the formulation buffer will minimize sticking if regular plastics are used.
- Avoid diluting lentivirus into low salt solution. Some lentiviruses aggregate in low salt solution, which will be non-infectious.

## Introduction of Lentivirus

---

Lentivirus (lente-, Latin for "slow") is a genus of retroviruses, causing chronic and deadly diseases by long incubation periods in human or other mammalian species [1]. To date, 5 serogroups have been recognized according to the vertebrate hosts they are associated with (primates, sheep and goats, horses, domestic cats, and cattle). Among them, the primate lentiviruses are distinguished by the utilization of CD4 surface protein as a receptor and the absence of dUTPase [2]. Derived from HIV-1, lentiviruses can integrate a significant amount of viral cDNA into the host genome, mediate stable and long-term transgene expression, and efficiently infect dividing cells and nondividing cells, which makes lentivirus an attractive gene delivery vehicle in most cell types [3].

Considering the key safety concerns during the use of HIV-derived lentivirus vectors, recombinant lentivirus has been designed and widely used for gene delivery in most cell types. As a research tool used to introduce a gene product into *in vitro* systems or animal models, lentiviral vector has been put into large-scale efforts to down-regulate or up-regulate gene expression in high-throughput formats, allowing researchers to examine the necessity and effects of transgenes in disease model systems, which is an indispensable for the discovery of novel transgenic drugs.

Nowadays, several generations of lentivirus packaging system are developed, in which the second-generation lentivirus vector and the third-generation lentivirus vector are the two most popular ones. The current method of the recombinant lentivirus production in Genemedi is based on three plasmids co-transfection system, involving the co-transfection of 3 plasmids (lentivirus series plasmid containing gene of interest (GOI) pLv-GOI, envelope expressing plasmid pMD2G and packaging plasmid pSPAX2) into 293T cells to generate lentivirus vectors.

## Protocol Overview

---

A schematic overview of recombinant lentivirus production is shown in Figure 1. The first step is to clone the gene of interest (GOI) into an appropriate LTR/MCS containing vectors.

The recombinant expression plasmid is co-transfected into the 293T cells with envelope expressing plasmid pMD2G and packaging plasmid pSPAX2, which together supply all of the trans-acting factors required for lentivirus replication and packaging in the 293T cells. Recombinant lentivirus particles are prepared from infected 293T cells and may then be used to infect a variety of mammalian cells.

Upon infection of the host cell, virus genome ssRNA should be converted into double-stranded DNA in order for

gene expression and virus replication. Together with other viral proteins, the newly synthesized DNA constitutes an integration-competent nucleoprotein complex, migrating into host cell nucleus and mediating integration of viral DNA into host chromatin. Integrated viral DNA, named as provirus, becomes part of host genome and serves as a transcription template for the synthesis of viral mRNA and genomic RNA. Following the synthesis of viral genomes and proteins, the viral components are assembled together to produce new virions, the virus particles then bud out of host cell and undergo a maturation step to generate infectious lentivirus.




Figure 1. Lentivirus packaging experiment flow chart.

## Experimental Materials

---

### Virus Packaging System

A three-plasmid system is used for packaging recombinant lentivirus (rLv) in this handbook, which includes a lentivirus series vector (pLv) that can be cloned into engineering sequences for gene overexpression, RNA interference and CRISPR/Cas9 gene knockouts, an envelope expressing plasmid pMD2G and a packaging plasmid pSPAX2. For more information regarding how to choose the right lentivirus vector for different experimental purpose, please consult our technical support.

### Bacterium Strain

E.coli strain DH5α is used for amplification of vectors.

## **Packaging Cell Line**

293T is the virus packaging cell line that can facilitate initial production, titer detection of lentivirus. It is an epithelial-like cell line required for lentivirus replication, and grows into a monolayer when confluent.

The complete growth medium of 293T is Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% Fetal Bovine Serum (FBS) and 1% Penicillin-Streptomycin (Pen-Strep). For a continuous culture, cells should not exceed 70% confluence to maintain proper characteristics. Usually, starting from cell passage number one, optimal results can be obtained within 30 passages. Once reached, it is best to start a new culture from another frozen stock in case of any unexpected mutations and unhealthy growth. Therefore, banking your own 293T frozen stocks is very important to ensure experimental integrity and continuity. Freezing cells at the logarithmic phase will improve post-thaw viability.

Note: To maintain cells in a healthier condition and improve production efficiency of lentivirus, it is recommended to use our Genemedi anti-mycoplasma reagent CurePlasma™.

## **Other Materials and Reagents**

Gene of interest  
LB broth  
Agar and Agarose  
Kanamycin  
Ampicillin  
70 and 100% ethanol  
Sterile PBS  
Chlorine bleach  
DNA gel apparatus and power supplies  
Class II Biosafety Cabinet  
37°C orbital shaker  
37°C bacteria incubator  
37°C, 5% CO<sub>2</sub> incubator  
15- and 50-ml conical tubes  
25- and 75-cm<sup>2</sup> tissue culture flasks  
Ultracentrifuge (Beckman) or equivalent with SW28 rotor  
Low-speed swinging-bucket centrifuge  
Microcentrifuge  
Centrifuge tube (thick-wall polycarbonate tube with cap)

## **Packaging and Concentration of Lentivirus**

---

### **Vector Construction of Lentivirus**

Before lentivirus packaging, gene of interest should be constructed into lentivirus vector. Genemedi has plenty of premade lentivirus vector goods carrying some genetic tools in stock, such as lentivirus-LC3 autophagy flux detection biosensors, etc.

**Note:**

In order to construct vectors quickly and efficiently, it is strongly recommended to use Genemedi- ClonEasy™ One Step Cloning Kit (Cat. GM-GC-01/02/03) .

**Transfection of Virus Plasmids into 293T Packaging Cells**

- a. Propagate 293T cells in DMEM with 10% FBS and 1% pen/strep. The day before transfection, plate the cells in a 10cm dish such that the cells reach 70-80% confluency the next day. On the day of transfection, set up the 3-plasmid co-transfection as table 1.
- b. DMEM needs to be preheated to 37°C with water bath. LipoGene™ transfection reagent needs to be warmed up to room temperature before use, and mix gently before use. Replace the transfection medium of 10cm dish with fresh medium 6 hours after transfection.
- c. To prepare viral plasmids for each reaction using a 60-mm dish:

**Table 1. Plasmid and transfection reagent required for transfection.**

Component	Amount
pLv-GOI vector	10µg
pMD2G	5µg
pSPAX2	10µg
LipoGene™	100µl

- d. Mix plasmids with transfection reagent in DMEM and add drop-wise to pre-seeded 293T cells. Incubate in 37°C, 5% CO2 and refresh with complete culture medium in 6 hours.

**Note:**

- 1. A detailed protocol of the transfection reagent can be referred to Genemedi LipoGene™ Transfection Reagent User Manual.
- 2. Cells should be in a healthy growth state for use prior to transfection.

**Harvest Virus**

Collect the supernatant containing lentivirus particles 48 hours and 72 hours later after transfection, respectively. Replace fresh DMEM culture medium after the collection of supernatant at 48h.

**Virus Purification**

- a. After collecting virus twice, discard the transfected 293T cells and filter the collected supernatant with 0.45µM filter membrane to an ultracentrifuge tube.
- b. Centrifuge at 72000g for 2 hours at 4°C.
- c. Then discard the supernatants and resuspend the lentivirus deposition with 500 µ l fresh medium and keep at -80°C or in liquid nitrogen for long time storage.

## Titration of Purified lentivirus

---

Determine lentivirus titer with fluorescent microscopy.

- a. Seed 293T  $1 \times 10^4$  cells/well in a 96-well plate one day in advance.
- b. Perform gradient dilution of the lentiviral particles to 1:10, 1:100, 1:1000,  $1:10^4$ ,  $1:10^5$ ,  $1:10^6$  in 100ul final volume in culture medium.
- c. Total 100 $\mu$ l viral particle mixture should be added to each well with at least 3 replicates per virus.
- d. Two days post infection, count the fluorescent positive cells using fluorescent microscopy and select the dilution factor with a proper fluorescent positive proportion (10%-30% positive cells/well). Count the triplicates and average the number of positive cells.
- e. Estimate the lentivirus titer using the following formulation: Viral titer (TU/ml) = number of fluorescent positive cells  $\times 10 \times$  dilution.

## Transduction of Target Cells

---

This protocol is for the stable cell line construction based on puromycin selection.

### **Note:**

*MOI: multiplicity of infection, is the number of viral particles to infect one cell. An optimization test of MOI is strongly recommended as the real MOI to certain cells may be affected by the operations and methods of dealing with viruses in different labs.*

### **Cell Preparation**

Plate robust target cells into 24-well plates at a density of  $1 \times 10^5$ /ml one day in advance.

### **Note:**

*The number of planted cells depends on the growth rate of the relevant cell line. 50% to 70% confluence should be reached on the following day.*

### **MOI Test of Lentivirus**

- a. Prepare the virus in 10-fold dilution gradient, and ensure the MOI is within a range of 3 to 1000.
- b. Day 0: Plate target cells in good condition at a density of  $1 \times 10^5$  /ml into 96-well plates, 100  $\mu$ l per well. Incubate at 37°C overnight.

- c. Day 1: Prepare virus in a six-MOI gradient, and dilute proper amount of virus suspension in complete culture medium of target cell to a final volume of 100  $\mu$ l (setting MOI = 3, 10, 30, 100, 300, 1000). Add diluted viruses to pre-seeded cells and incubate for 4 to 8 hours at 37°C, then refresh the medium to remove viruses.
- d. Day 3: Detect fluorescence with a microscope. Calculate MOI based on the ratio of fluorescent cells.

**Note:**

If the virus is not fluorescence-labeled, MOI can be determined by qPCR, WB, IF, IHC, etc.

**Table 2. The lentiviral MOIs of commonly used cell lines.**

Cell line	MOI range	Auxiliary infection reagent polybrene
<b>K562</b>	20~40	Need
<b>Jurkat</b>	50~80	No
<b>kasumi</b>	10~30	No
<b>NB4</b>	50~80	No
<b>U937</b>	20~40	Need
<b>THP-1</b>	50~80	Need
<b>GBC-SD</b>	30~50	No
<b>H929</b>	100~150	No
<b>H1299</b>	1~3	Need
<b>95D</b>	2~4	Need
<b>A549</b>	20~40	Need
<b>SPC-A-1</b>	100~150	Need
<b>7402</b>	10~15	Need
<b>Hep 3B</b>	10~30	Need
<b>Hep G2</b>	10~30	Need
<b>SMMC-7721</b>	10~30	Need
<b>Huh-7</b>	10~30	Need
<b>Hela</b>	10~30	Need
<b>HOS</b>	20~40	Need
<b>Hep-2</b>	10~30	Need
<b>HL-60</b>	>100	Need
<b>HT-29</b>	10~30	Need
<b>PKO</b>	2~4	Need
<b>SW480</b>	10~30	Need
<b>DLD-1</b>	10~30	Need
<b>SK-OV-3</b>	2~4	Need
<b>SHG-44</b>	10~30	Need
<b>U251</b>	1~3	Need
<b>U87</b>	1~3	Need
<b>293T</b>	1~3	Need
<b>HUVEC-2C</b>	10~30	Need
<b>PC-3</b>	20~40	Need
<b>MDA-MB-</b>	10~30	Need
<b>MCF-7</b>	20~40	No
<b>Tca8113</b>	10~30	Need

<b>RPE</b>	10~30	Need
<b>AGS</b>	100~150	Need
<b>BGC-823</b>	100~150	Need
<b>SGC-7901</b>	10~30	Need
<b>MKN-28</b>	20~40	Need
<b>MKN-45</b>	20~40	Need
<b>BxPc-3</b>	20~40	Need
<b>CFPAC-1</b>	50~80	Need
<b>Panc-1</b>	2~4	Need
<b>HEC-1-B</b>	2~4	Need
<b>NIH-3T3</b>	20~40	Need
<b>Raw264.7</b>	10~30	No
<b>CHO</b>	20~40	Need
<b>HSC-T6</b>	10~30	No
<b>C6</b>	>100	Need
<b>NRK</b>	10~30	Need

### **Transduction**

- a. Before infection, virus should be melted on ice gently and resuspended in culture medium.
- b. Prepare the virus in 10-fold dilution gradient, and ensure the MOI is within a range of 3 to 1000.
- c. Remove the preceding medium and add lentivirus-containing medium with 1/2 volume of normal culture volume.
- d. Culture for 4 hours at 37 °C , and supplement fresh medium to normal volume. The recommended medium volume of lentivirus infection is displayed in the following table 3.
- e. Refresh the culture medium 24 hours post infection.

**Table 3. The recommended medium volume during lentivirus infection.**

<b>Culture</b>	<b>Surface</b>	<b>Normal volume for cell</b>	<b>1/2 Volume for lentivirus</b>
96-well	0.3 cm <sup>2</sup>	100 µl	50 µl
24-well	2 cm <sup>2</sup>	500 µl	250 µl
12-well	4 cm <sup>2</sup>	1 ml	500 µl
6-well	10 cm <sup>2</sup>	2 ml	1 ml

### **Construction of Stable Transgenic Cell Lines**

48 hours post infection, change to fresh medium with puromycin. The recommended concentration of puromycin ranges from 1 to 10 µg/ml according to cell lines. Set the uninfected wild-type cells as control group and add equal volume and concentration of puromycin. Replace with fresh puromycin-containing medium every 2 or 3 days until the control group cells die out. Then choose one of the following steps according to experimental requirements.

- a. Non-selecting monoclonal cells

Passage the infected cells and select with puromycin constantly. Freeze the cell mixture in continuous three passages.



Considering the heterogeneity, we recommend selecting monoclonal cell for a confirmed phenotype.

b. **Selecting monoclonal cells**

Select at least five monoclonal cells after infection and puromycin selection, and propagate in puromycin-containing medium. Detect the expression of target genes using western blot or qPCR. Choose the stable cell line with proper expression level of target genes to passage three generations and freeze the stable cell line.

***Notes for infection of special cell lines:***

1. Suspension cells

We recommend using flat fillet centrifuging transfection to infect suspension cells or semi-suspension cells. Add virus suspension into cell culture dish, sealing tightly, and centrifuge at low speed of 200g for 1 hour in the flat fillet centrifuge. Place cells in cell culture incubator after centrifuging transfection. If the flat fillet centrifuge is inaccessible, you can suspend the cells and transfer cells into centrifuge tubes, followed by low-speed centrifuge, and discard the most of supernatant. Add virus suspension into the tubes, resuspending cells, place it at room temperature for 15 min (no more than 30 min), and transfer the cells and virus suspension into plate to culture. Replace with fresh culture medium the next day.

2. Cells difficult to infect

For cells difficult to infect, like DC cells, we recommend repeated infections. Replace with fresh virus suspension 24 hours after the first infection. Repeated infections can increase the infection efficiency markedly.

3. Non-dividing primary cells

We recommend high-titer adenovirus to infect these cells like BMSC.

## References

---

1. Salgado CD and JM Kilby. (2009). Retroviruses and other latent viruses: the deadliest of pathogens are not necessarily the best candidates for bioterrorism. J S C Med Assoc 105:104-6.
2. Piguet V, O Schwartz, S Le Gall and D Trono. (1999). The downregulation of CD4 and MHC-I by primate lentiviruses: a paradigm for the modulation of cell surface receptors. Immunol Rev 168:51-63.
3. Cockrell AS and T Kafri. (2007). Gene delivery by lentivirus vectors. Mol Biotechnol 36:184-204.

## Contact Information

---

Genemedi Biotech. Inc.

For more information about lentivirus, please visit: [www.genemedi.net/i/lentivirus-packaging](http://www.genemedi.net/i/lentivirus-packaging)

For more information about Genemedi products and to download manuals in PDF format, please visit our web site:

[www.genemedi.net](http://www.genemedi.net)

For additional information or technical assistance, please call or email us

Worldwide: [+86-21-50478399](tel:+86-21-50478399)

Fax: [+86-21-50478399](tel:+86-21-50478399)

E-mail: [support@genemedi.net](mailto:support@genemedi.net)

© 2018 Genemedi Biotech. Inc. All rights reserved.